

CELL CULTURE MEDIA FOR MAMMALIAN CELLS

FIELD OF THE INVENTION

The present invention relates generally to cell culture media for mammalian cells. In particular, the invention relates to cell culture media that allow long-term expansion and maintenance of a cell population of mammalian hepatocytes, hepatocyte-derived cell lines, hepatocyte-derived malignant cells, and other cells.

BACKGROUND OF THE INVENTION

It is well-known that specific cell lines can be grown in vitro in optimally formulated culture or nutrient media. Examples of some of the culture media developed for special purposes are: RPMI Media 1640 medium for growth of human B-lymphoid cells and malignant cells, Changs medium for growth of amniotic fluid cells, Medium 199 for growth of mouse fibroblast cells, Minimal Essential Medium (MEM) medium, a "minimal" medium for growth of attached mammalian cells, and Leibovitz medium for growth in absence of CO₂. Such various media are distinguished from one another in that they contain critically different components in precise amino acids, vitamins, organic salts, trace elements, and other organic compounds which promote the maximum growth of the cultured cells.

For the growth of mammalian cells chemically defined media are usually supplemented with various sera, preferably fetal calf or newborn calf serum, and other incompletely defined growth factors. A major drawback to serum, however, is that its constituents may vary widely, thereby introducing undefined biological components into the nutrient medium which contributes to the variability

of biochemical and cellular events. Additionally, serum is expensive and may result in critical immune reactions in patients if the cells are used for clinical purposes.

5 The present invention is primarily directed to culturing mammalian hepatocyte cells using chemically defined media that allow long-term expansion of the cell population. The term "chemically defined media" is used in tissue culture to refer to culture media of known
10 chemical composition, both quantitatively and qualitatively, as contrasted with media which contain natural products such as animal serum.

 Liver regeneration is achieved primarily by cell division of mature adult hepatocytes as reported by
15 Grisham, J.W., et al., Cancer Res. 22:842 (1962), the disclosure of which is incorporated herein by reference. These cells, or a fraction thereof, have a high capacity for clonal growth, as shown by hepatocyte transplantation experiments in ectopic sites (Jirtle, R.L., et al., Cancer
20 Res. 42:3000 (1982), the disclosure of which is incorporated herein by reference) and in transgenic mouse models (Rhim, J.A., et al., Science 263:1149 (1994), the disclosure of which is incorporated herein by reference). It has been shown in several studies, however, that when
25 liver is stimulated to regenerate while proliferation of mature hepatocytes is suppressed, facultative stem cells emerge and proliferate. See, for example, Thorgeirsson, S.S., et al., Proc. Soc. Exp. Biol. Med. 204:253 (1993), the disclosure of which is incorporated herein by
30 reference. Such cells, sometimes referred to as "oval cells," can mature into hepatocytes in defined animal models or ductular structures composed of cells ("ductular hepatocytes") with mixed hepatocyte and bile duct epithelial markers. See, Gerber, M.A., et al., Amer. J.
35 Path. 110:70 (1983) and Vandersteenhoven, A.M., et al., Arch. Pathol. Lab. Med. 114:403 (1991), the disclosures of

which are incorporated herein by reference. Little is known, however, about their origin and about the controls that regulate their phenotypic transitions to hepatocytes or ductular cells.

Despite the high capacity of hepatocytes to proliferate in vivo, directly or via facultative stem cell growth, the conditions that determine their growth potential and their phenotypic transitions are not thoroughly understood because of only limited success in hepatocyte growth in primary culture. It is typically the case that hepatocytes in primary culture under the influence of primary mitogens enter into one or two divisions and then the cells degenerate and die. Heretofore various investigators have failed to develop a medium that allows hepatocytes to both proliferate and survive.

For example, Berry, N.M., et al., J. Cell. Biol. 43:506 (1969), the disclosure of which is incorporated herein by reference, taught the collagenase perfusion technique which allows liver tissue to dissociate into its component cellular elements, based on size. Later, Bissell, D.M., et al., J. Cell. Biol. 59(3):722 (1973) and Bonney, R.J., et al., In Vitro 2:399 (1974), the disclosures of which are incorporated herein by reference, described the first methods for culturing isolated hepatocytes which perhaps survived for one or two days. Long term culture of hepatocytes on collagen gels for a maximum of 7 to 10 days was reported by Michalopoulos, G., et al., Exp. Cell. Res. 94(1):70 (1975), the disclosure of which is incorporated herein by reference. The common characteristic of all of the above-referenced systems is that the hepatocytes in those systems were maintained in culture without there being any evidence of cell

proliferation. These systems were, instead, only maintenance cultures of non-proliferating cells for a brief period.

5 The first successful attempt to initiate DNA synthesis in hepatocytes used the then newly-discovered epidermal growth factor (EGF) as reported by Richman, R.A., et al., Proc. Nat. Acad. Sci. USA 73:3589 (1976), the disclosure of which is incorporated herein by
10 reference. Over the ensuing years several other groups of researchers used EGF as a mitogen for hepatocytes and reported on the mitogenic effects of EGF and their modulation by other factors such as, for example, matrices such as collagen Type I, zinc, and proline.

15 Hepatocyte growth factor, also known as scatter factor (hereinafter referred to as "HGF" or "HGF/SF") was discovered, cloned and ultimately sequenced by the late 1980's. See Michalopoulos, G., et al., Federation Proceedings 42:1023 (1983); Michalopoulos, G., et al.,
20 AACR Proceedings 24:58 (1983); Michalopoulos, G., et al., Cancer Res. 44(10):4414 (1984); and Miyazawa, K., et al., Biochem. Biophys. Res. Commun. 163:967 (1989), the disclosures of which are incorporated herein by reference. HGF/SF was found to be a mitogen for many hepatocytes as
25 well as for epithelial cells. HGF/SF's importance for the liver is due to the fact that it is the trigger for liver regeneration through an endocrine mechanism.

 Recently, several studies have shown that HGF/SF, epidermal growth factor ("EGF"), and transforming
30 growth factor α ("TGF α ") are the primary mitogens for hepatocytes in culture by stimulating limited hepatocyte DNA synthesis in chemically defined media. See, for example, Michalopoulos, G.K., Fed. Am. Soc. Exp. Biochem J. 4:176 (1990), the disclosure of which is
35 incorporated herein by reference. These growth factors

were later found to additionally play a role in vivo in liver regeneration after partial hepatectomy. Injection of HGF/SF, TGF α , or EGF in rats induces DNA synthesis in hepatocytes. See, for example, Liu, M.L., et al., Hepatology 19:1521 (1994), the disclosure of which is incorporated herein by reference.

In all of these systems, however, it was reported that hepatocytes entered into DNA synthesis and mitosis for only a limited time, typically 1-3 days. After 1-2 rounds of DNA synthesis and cell division, the cultures degenerated and all the cells were dead in about 7-10 days. Until now, there has also been no documented expansion of the number of hepatocytes in cell culture by adding either EGF or HGF alone, or in combination. The cell replication in cultures containing these growth factors is instead self-limited and the number of hepatocytes that die exceeds the number of hepatocytes that are newly generated. The cell replication in cultures containing other hepatocyte mitogens such as transforming growth factors, such as TGF α , and acidic fibroblast growth factor is similarly self-limited.

More recently, Mitaka, T., et al., Hepatology 13(1):21 (1991); Mitaka, T., et al., Hepatology 16(2):440 (1992); Mitaka, T., et al., Virchows Arch. B Cell Pathol. Incl. Mol. Pathol. 62:329 (1992); and Mitaka, T., et al., Cancer Res. 53:3145 (1993), the disclosures of which have been incorporated herein by reference, have reported that adding nicotinamide, dexamethasone and EGF to a conventional culture medium resulted in the formation of colonies of small hepatocytes arising in a dense culture of standard size parenchymal hepatocytes. In a further study, Mitaka, T., et al., J. Cell. Physiol. 157:461 (1993), the disclosure of which is incorporated herein by reference, have reported that the numbers of colonies induced by the combinations of EGF + HGF, EGF + TGF- α , and

HGF + TGF- α were not different from those of colonies induced by each mitogen alone. In these studies, however, there was no significant expansion of the total cell population, no evidence of clonal growth, and there was loss of differentiation.

Until now there has been no chemically defined medium, supplemented or not, which is able to support long term proliferation, differentiation, and viability of hepatocytes. While for many purposes the use of an undefined supplement is satisfactory, in cases where studies are made of growth, metabolism, and/or differentiation of cells in culture, it is most desirable to have a supplement that is defined. The introduction of undefined components to a cell culture can contribute to variability, unpredictability, and contamination in study results and applications of cell cultures. The use of defined media is particularly important and advantageous in areas of drug metabolism, artificial organ development, cell transplantation, gene therapy, and basic investigational cell studies.

The above-described limited capacity of hepatocytes to proliferate in primary culture has hindered long-term studies or uses that required long-term viability or proliferation. Applications of hepatocyte cultures to cellular transplantation and gene therapy have thus been hindered. Consequently, there remains a need for a chemically defined medium that will allow hepatocytes to proliferate and survive long term. Among the potential applications for such a medium and the hepatocytes and other cells so cultured are gene therapy, bioartificial organs, cell transplants, drug production, and drug and chemical testing.

As stated above, the current state of the art does not provide a hepatocyte culture system in which hepatocytes expand as a cell population by sustained proliferation, and there is a need for such a system. The present invention provides a fully defined culture medium which allows sustained proliferation and long-term expansion of hepatocytes.

SUMMARY OF THE INVENTION

According to the present invention, a new chemically defined cell culture medium is provided. This medium supports sustained clonal growth of primary hepatocytes and hepatocyte cell lines, genetically transformed hepatocytes, and hepatocytes obtained from neoplastic sources, resulting in expansion of the cell population. This medium further allows complete differentiation of metabolic, structural, and secretory functions of the cells grown therein. Under these conditions, hepatocytes undergo multiple proliferative cycles. Once confluency is reached, or in the presence of specific matrix components, nutrients, and/or growth factors, these proliferating cells stop dividing and maintain a mature hepatocyte phenotype for many months or longer.

Accordingly it is a primary object of the present invention to provide a culture medium for sustained proliferation and viability of hepatocytes.

Another object of the present invention to provide a culture medium for sustained differentiation and viability of hepatocytes.

Yet another object of the present invention is to provide a culture medium for sustained proliferation of hepatocytes that revert to complete differentiation as growth ceases.

Another object of the present invention is to provide a culture medium for long-term expansion of hepatocytes that contains no serum such that the medium is fully defined.

Yet another object of the present invention is to provide a culture medium for sustained proliferation, differentiation, and viability of hepatocytes on a variety of matrix substrates.

These and other objects of the present invention are achieved by one or more of the following embodiments.

In one aspect, the invention features a chemically defined HBM culture medium for maintenance, differentiation, and long-term growth of mammalian hepatocytes, comprising:

(a) a synthetic stock basal medium designed for mammalian cell culture; and

(b) a hepatocyte cell growth promoting amount of components selected from among nicotinamide, amino acids, transferrin, hormones, dexamethasone, trace metals, and simple carbohydrate selected from the group consisting of D-glucose and D-galactose and any combination thereof.

In a preferred embodiment the invention features HBM culture medium further comprising buffer, antibiotics, and albumin.

In another aspect, the invention features a mammalian cell culture medium comprising the composition of HGM as defined in Tables I and II, wherein the stock
5 basal media of Table I comprises a blended DMEM such that the final concentration of D-glucose is preferably about 2.0 g/L and the amount of D-galactose is preferably about 2.0 g/L.

Other features and advantages of the invention
10 will be apparent from the following description of the preferred embodiment, and from the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A-1C are graphs showing results of studies in which rat hepatocytes were cultured in HBM
15 medium as described herein supplemented with growth factors, as indicated. All points represent the mean and standard error of at least three separate cultures.

Figure 1A shows the percent labeled nuclei (BRdU) of proliferating cell cultures at different times
20 after hepatocyte isolation. The cells were grown in HBM supplemented with HGF/SF and EGF.

Figure 1B shows the incorporation of [³H] thymidine (disintegrations per minute) into DNA in cultures at different times after hepatocyte isolation.
25 Cells were exposed to HGF/SF in HBM (◇); EGF in HBM (○); HGF/SF + EGF in HBM (Δ); and a control, HBM alone (□).

Figure 1C shows the amount of DNA per plate at different days of cells grown in HBM medium alone (control) (■); HGF/SF in HBM (Δ); EGF in HBM (▽); and
30 HGF/SF + EGF in HBM (●).

Figure 2 is a graph showing the day 15 DNA per plate of rat hepatocytes grown in HBM with the indicated growth factors. Cells represented by control (day t=0) and day 15 were grown in HBM without any growth factors.

Figure 3 is a graph showing the amount of DNA synthesis per plate ($\mu\text{g/culture}$) of rat hepatocytes at day 15 in HBM supplemented with HGF/SF and EGF grown on different matrices. "CC" means collagen coated; "ECL" is a commercial matrix; "m COLL IV" means mouse collagen IV; "CC VITROGEN" means bovine skin collagen type I; "POLY D LYS" means poly D-Lysine; "m LAMININ" means mouse laminin; "h FIBRONECT" means human fibronectin; and "m COL I" means mouse collagen I.

Figures 4A-4G are photographs of rat hepatocytes cultured in HBM as described herein under various conditions.

Figure 4A shows hepatocytes in HBM medium with HGF/SF and EGF at day 1 after isolation showing typical subconfluent non-proliferating hepatocytes.

Figure 4B shows hepatocytes at day 4 in HBM media induced with HGF/SF showing typical scattered morphology.

Figure 4C shows hepatocyte cultures having reached confluency at day 15 showing typical morphology.

Figures 4D and 4E are election photomicrographs of the cells in Figures 4A and 4C, respectively.

Figures 4F and 4G are photomicrographs of stained hepatocytes that were transfected at day 3 with a lac-Z-containing replication defective retrovirus and stained for β -galactosidase expression as described below.

and then photographed at day 1 (Fig. 4F) and day 10 (Fig. 4G) after transfection. The cells were cultured in HBM medium supplemented with HGF/SF and EGF as described below.

Figures 5A and 5B are photographs of Northern blots showing expression of specific genes at different days in rat hepatocyte cultures maintained in HBM in the presence of HGF/SF and EGF as described herein. GAPDH expression and the intensity of the 28S RNA after ethidium bromide staining were used as internal controls.

Figures 6A-6F are photographs of proliferating rat hepatocytes cultured in HBM medium in the presence of HGF/SF and EGF as described herein and Northern blots thereof.

Figure 6A is a phase contrast photomicrograph of cultures of proliferating hepatocytes overlaid with Matrigel at day 8 and photographed at day 18. The granular cytoplasm and the appearance of typical bile canaliculi appear as bright lines between cells.

Figures 6B and 6C are low and high power electron photomicrographs, respectively, of cells at day 18 in culture, 10 days after overlay with Matrigel. Typical features of hepatocyte cytoplasm are shown such as lamellae of endoplasmic reticulum wrapping around mitochondria, microbodies with crystalline center, bile canaliculi ("c") (Figure 6B), abundant mitochondria ("M"), and glycogen ("G") (both in Figure 6C).

Figure 6D is a photograph of a Northern blot showing increased expression of albumin mRNA after addition of Matrigel. Albumin mRNA is expressed in control cultures, immediately after isolation from liver by collagenase perfusion and before culture. Expression

is minimal at day 8 in culture. At days 3 and 7 after Matrigel addition (lanes marked with "+"), there was an increase in expression of albumin mRNA. GAPDH expression was used as an internal control.

Figure 6E is a photograph of a Northern blot showing induction of cytochrome IIB1 mRNA in cultures treated with Matrigel at day 8 and exposed to phenobarbital ("PB") 2 days later (day 10 of culture). Cells were harvested at day 15 of culture. GAPDH expression was used as an internal control for mRNA loading.

Figure 6F is a photograph of a Northern blot of cells cultured additionally with Matrigel (added at day 8 in culture). Cytokeratin 19, a bile duct marker expressed by the proliferating hepatocytes, is suppressed by addition of Matrigel and is not suppressed in control cultures not receiving Matrigel. 28S rRNA stained by ethidium bromide was used as an internal control.

Figures 7A-7E show the results of studies showing formation of ductular/acinar structures in cultures of rat hepatocytes kept from the beginning of culture between two type I collagen gel layers in HBM in the presence of HGF/SF. The Figures 7A-7D photographs are of 15 day cultures.

Figure 7A is a phase contrast photomicrograph (100X) showing the appearance of the ductular structures surrounded by collagen fibrils.

Figure 7B is a photomicrograph (100X) of paraffin sections of the Figure 7A type I collagen gels stained with hematoxylin and eosin.

Figures 7C and 7D are electron micrographs of cells surrounding the same lumen but in different locations around the lumen of one of the ductular structures seen in Figure 7B. Figure 7C shows cells with a morphology similar to bile duct epithelium, with long parallel contacts joined by many desmosomes and with abundant keratin intermediate filaments. Figure 7D shows cells resembling more of the hepatocyte phenotype, with rough endoplasmic reticulum and mitochondria, densely stained secondary lysosomes and fewer filaments.

Figure 7E is a photograph of a Northern blot showing that expression of cytokeratins 18 and 19 increases in cultures with ductular/acinar structures whereas albumin is only slightly expressed.

Figure 8 shows photographs of stained human hepatocyte cultures grown in HBM supplemented with HGF/SF and EGF as described herein taken at day 1, 3, 5, 7, 10, 12, and 19.

20

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a hepatocyte basal medium, designated HBM herein, which allows long-term expansion, differentiation and survival of the cell population of hepatocytes, hepatocyte-derived cell lines such as HepG2, hepatic fetal epithelial cells, and hepatic primary hepatocarcinoma cells in vitro. Additionally, the media of the present invention can be used for culturing pancreatic islet cells, renal tubular cells, Ito cells, small intestine epithelial cells, and a variety of cell lines, including MRC5, CaCo, and 3T3 cells, as well as others. The HBM of the present invention maintains metabolic pathways and synthesis functions, i.e., differentiation of mature adult hepatocytes. Additionally, the HBM of the present invention, when

combined with specific mitogens and extracellular matrix compositions, hepatocytes can be driven to transdifferentiate into bile duct-like structures. The HBM of the present invention can be used for culture of mammalian hepatocytes, and other cells, including, but not limited to those of human, rat, dog, pig, mouse, and baboon origin.

The HBM medium comprises appropriate levels of essential and non-essential amino acids and bulk ions and trace elements, buffers, vitamins, carbohydrates, lipids, proteins, and hormones to function as a nutrient medium for in vitro mammalian cell culture.

In its broadest aspect the invention features a chemically defined basal media, HBM, that by itself allows for long-term survival, differentiation, and growth of mammalian hepatocytes and other cells. Additionally, in the presence of growth factors such as HGF/SF, EGF, or TGF α , as well as other mitogens, cells growing in the supplemented HBM have a more rapid population expansion and clonal growth. As further demonstrated below, the HBM of the present invention when supplemented with HGF/SF causes formation of bile duct-like structures in conjunction with certain matrix constructs. As stated above, however, such growth factors are not required for specific differentiation patterns in primary cultures, but do accelerate growth and population expansion if this is desired for a specific purpose.

The HBM medium of the present invention comprises a chemically defined stock basal medium (hereinafter referred to as "SM") designed for mammalian cell culture. The stock basal medium can be preferably constructed using Dulbecco's Modified Eagle Medium ("DMEM") as one ingredient, but the present invention is not to be so limited as long as the formulations are within the guidelines set forth herein. Examples of other

defined basal medias which may be used in accordance with the present invention include, but are not limited to: Basal Media Eagle (BME), DMEM/F-12 (1:1 DMEM and F-12 vol:vol); Medium 199; F-12 (Ham) Nutrient Mixture; F-10 (Ham) Nutrient Mixture; Minimal Essential Media (MEM), Williams' Media E; and RPMI 1640, all of which are available from Gibco - BRL/Life Technologies, Inc., Gaithersburg, MD, among others. Several versions of many of these media are available, and those that are particularly useful to construct HBM include, but are not limited to: DMEM 11966, DMEM 10314, MEM 11095, Williams' Media E 12251, Ham F12 11059, MEM-alpha 12561, and Medium-199 11151 (all available from Gibco-BRL/Life Technologies (1995-1996 catalog)). Therefore, for example, if L-arginine and/or D-glucose are already in the stock basal media in the necessary amounts, then little or no additional ingredient will have to be added as a supplement.

Depending on the particular composition of the stock basal medium, the SM is then supplemented, as described more fully herein, with D-glucose and/or D-galactose, nicotinamide, other micronutrients comprising amino acids and trace metals not already present in the SM such as L-proline, L-glutamine, L-arginine, L-ornithine, zinc, manganese, copper, and selenium, purified transferrin to which is bound elemental iron or apo-transferrin in combination with iron gluconate, hormones such as dexamethasone and insulin and a pH buffer such as HEPES (N-[2-hydroxyethyl]piperazine-N-[2-ethanesulfonic acid]). Antibiotics such as penicillin and streptomycin, a pH indicator, albumin and/or dextran, essential fatty acids, alternative buffers, vitamins, osmotic agents, and other forms of trace metals may also be optionally added. Typically, a basal medium will have a pH in the range of 6.5-8.2, preferably 7.0-7.7, and most

preferably 7.2-7.5. Phenol red is a typical indicator added to aid in the control of pH. The SM and supplements thereto comprise the HBM medium of the present invention.

5 The HBM medium may then be optionally supplemented with one or more growth factors such as, for example, HGF/SF, EGF and TGF α if accelerated growth is desired.

10 Simple carbohydrates D-glucose and/or D-galactose are added to the stock basal media to comprise HBM. If both D-glucose and D-galactose are used, the sum of their total concentrations is preferably 8.0 g/L or less but greater than .01 g/L, taking into account the amount of D-glucose present in the stock basal media, if
15 any. When only one of D-glucose or D-galactose is used, the concentration is preferably 5.0-0.1 g/L. For example, in the Example herein, the blended DMEM stock basal medium that was used contained 2.0 g/L D-glucose and no further D-glucose was added. 2.0 g/L of D-galactose was then
20 added as a supplement.

 Nicotinamide, another HBM component, has been shown to maintain hepatocyte differentiation, enhance expression of cytochrome P450, and prolong hepatocyte survival in conventional culture to 10-14 days. See,
25 Rosenberg, M.R., et al., In Vitro 18: 775 (1982) and Inoue, C., et al., Biol. Chem. 264(9): 4747 (1989), the disclosures of which are incorporated herein by reference.

 Transferrin is an iron binding protein that interacts with a transferrin receptor on the cell
30 membrane. It serves to both chelate and transport iron ions. The transferrin preferably used in the present invention is either holo-transferrin 30% saturated with iron or is completely unsaturated (apo-transferrin) and is combined with iron gluconate.

Dexamethasone, a synthetic corticosteroid, has been shown to enhance EGF-induced DNA synthesis as reported by Sand, T.-F., et al., Acta. Endocrinol. 109:369 (1985), the disclosure of which is incorporated herein by reference. As used in the present invention, dexamethasone can be any cortisol derivative such as, prednisone, prednisilone, cortisol, hydrocortisone, and other derivatives.

Insulin and insulin-like growth factors are required for glucose uptake, amino acid transport, and maintenance of multiple intermediary metabolic pathways. These effects help to maintain differentiation and support proliferation.

The inclusion of L-arginine in HBM, in the stock basal medium, or supplemental thereto, appears to be important because hepatocytes in culture tend to lose their capacity to synthesize arginine through the urea cycle. In the absence of L-arginine, hepatocytes in culture cannot live for very long because they become incapable of synthesizing L-arginine, thereby blocking their protein synthesis. The use of D-galactose in addition to D-glucose is advantageous for HBM because the combination gives the greatest growth potential over either substance alone.

HGF/SF, EGF, and TGF α , as discussed above, are mitogens which as shown below exhibit an enhanced proliferative effect when used together or alone in accordance with the present invention. The foregoing list of mitogens that can be used to supplement the HBM of the present invention is not exhaustive. It is to be noted, however, that these mitogens are not required for survival or differentiation of hepatocytes when grown in HBM. Hepatocytes grown in HBM will proliferate at a slower rate than if mitogens are added.

The insulin, EGF, HGF, and TGF α used in the presently claimed media can either be recombinantly produced, genetically engineered, or purified from natural sources. The species can be, for example, human, bovine, equine, murine, porcine, or rat.

Several preferred stock basal media - DMEM 11966, DMEM 10314, MEM 11095, and Williams' Medium E 12251 contain the following components per 1000 ml of sterile deionized water as shown below in Table I:

TABLE I
Compositions of Stock Basal Media (mg/L)

	COMPONENT	DMEM 11966	DMEM 10314	MEM 11095	WILLIAMS' MEDIUM E 12251
	<u>INORGANIC SALTS:</u>				
	CaCl ₂ (anhyd.)	200.00	200.00	200.00	200.00
	CuSO ₄ · 5H ₂ O	--	--	--	0.0001
	Fe (NO ₃) ₃ · 9H ₂ O	0.10	0.10	--	0.0001
20	KCl	400.00	400.00	400.00	400.00
	MnCl ₂ · 4H ₂ O	--	--	--	0.0001
	MgSO ₄ · (anhyd.)	97.67	97.67	97.67	97.67
	NaCl	6,400.00	6,400.00	6,800.00	6,800.00
	NaHCO ₃	3,700.00	3,700.00	2,200.00	2,200.00
25	NaH ₂ PO ₄ · H ₂ O	125.00	125.00	140.00	--
	NaH ₂ PO ₄	--	--	--	140.00
	ZnSO ₄ · 7H ₂ O	--	--	--	0.0002

OTHER COMPONENTS:

	D-Glucose	--	4,500.00	1,000.00	2,000.00
	Glutathione	--	--	--	0.05
5	Methyl Linoleate	--	--	--	0.03
	Phenol red	15.00	15.00	10.00	10.00
	Sodium Pyruvate	--	--	--	25.00

AMINO ACIDS:

	L-Alanine	--	--	--	90.00
10	L-Arginine·HCl	84.00	84.00	126.00	--
	L-Arginine	--	--	--	50.00
	L-Asparagine·H ₂ O	--	--	--	20.00
	L-Aspartic Acid	--	--	--	30.00
	L-Cysteine	--	--	--	40.00
15	L-Cystine	--	48.00	--	--
	L-Cystine·2HCL	63.00	63.00	31.00	26.10
	L-Glutamine	584.00	584.00	292.00	--
	L-Glutamic Acid	--	--	--	50.00
	Glycine	30.00	30.00	--	50.00
20	L-Histidine·HCl·H ₂ O	42.00	42.00	42.00	--
	L-Histidine	--	--	--	15.00
	L-Isoleucine	105.00	105.00	52.00	50.00
	L-Leucine	105.00	105.00	52.00	75.00
	L-Lysine·HCl	146.00	146.00	73.00	87.50
25	L-Methionine	30.00	30.00	15.00	15.00
	L-Phenylalanine	66.00	66.00	32.00	25.00
	L-Proline	--	--	--	30.00
	L-Serine	42.00	42.00	--	10.00
	L-Threonine	95.00	95.00	48.00	40.00
30	L-Tryptophan	16.00	16.00	10.00	10.00
	L-Tyrosine·2Na·2H ₂ O	104.00	104.00	52.00	50.70
	L-Valine	94.00	94.00	46.00	50.00

VITAMINS:

	Ascorbic Acid	--	--	--	2.00
	Biotin	--	--	--	0.50
5	D-Ca pantothenate	4.00	4.00	1.00	1.00
	Choline chloride	4.00	4.00	1.00	1.50
	Ergocalciferol	--	--	--	0.10
	Folic acid	4.00	4.00	1.00	1.00
	i-inositol	7.20	7.20	2.00	2.00
10	Menadione				
	Na Bisulfate	--	--	--	0.01
	Niacinamide	4.00	4.00	1.00	1.00
	Pyridoxine HCl	--	4.00	--	--
	Pyridoxal HCl	4.00	--	1.00	1.00
15	Riboflavin	0.40	0.40	0.10	0.10
	α -Tocopherol				
	Phosphate Disodium	--	--	--	0.01
	Thiamine HCl	4.00	4.00	1.00	1.00
	Vitamin A Acetate	--	--	--	0.10
20	Vitamine B ₁₂	--	--	--	0.20

In accordance with the preferred embodiment of present invention, the following supplemental components in Table II are added to the stock basal media of Table I in a total volume of 1000 ml. The preferred amounts listed coincide with those described below in the Example for which the basal stock medium was constructed from 445 ml of DMEM 10314 and 555 ml 11966 to provide a D-glucose concentration of 2.0 g/L such that no additional D-glucose was added. It is to be understood that when using other basal stock media the optimal amounts of these components that are added may vary.

TABLE II
Additions to SM for HBM

5	Preferred		Concentration Range
	Amount		
	units/L		units/L
	D-Galactose	2.0 g	0.01-5.0 g*
	D-Glucose	2.0 g**	0.01-5.0 g*
	Nicotinamide	610.0 mg	1-3050 mg
10	L-Proline	30.0 mg**	1-120 mg
	L-Arginine	84.0 mg**	1-150 mg
	L-Ornithine	100 mg	1-500 mg
	human-holo-Transferrin (30% Fe saturated)	5.0 mg	0.1-100 mg
15	h-Insulin	5.0 mg	greater than 10 ⁻¹¹ M
	Dexamethasone	1 x 10 ⁻⁷ M	10 ⁻¹² -10 ⁻³ M
	ZnCl ₂	.544 mg**	1-3000 µg
	MnSO ₄	0.025 mg**	1-250 µg
	ZnSO ₄ ·7H ₂ O	.750 mg**	1-3000 µg
20	CuSO ₄ ·5H ₂ O	0.20 mg**	1-1000 µg
	Selenium	5.0 µg	1-150 µg
	L-Glutamine*** (additional to SBM)	5.0 mM	2.0-10.0mM
	HEPES	20.0 mM	5-50 mM
25	* If both D-galactose and D-glucose are used, the upper limit is about 8.0 g/L or less, taking into consideration the amounts present in the particular stock basal medium used, if any. The lower limit for when both or only one of D-glucose or D-galactose is		
30	used is about 0.01 g/L or greater. For example, in the Example herein, the blended DMEM medium used contained 2.0 g/L D-glucose with no additional amount added, and 2.0 g/L D-galactose was added.		
	** If not already in SM stock basal medium.		
35	*** L-glutamine is a required nutrient in the present invention. Many stock basal medias are provided with L-glutamine included. However, it is well known by		

5 practitioners of the art of cell culture that
 L-glutamine will degrade by oxidation, such that all
 L-glutamine will be degraded over a period of a few
 weeks following manufacture. Therefore, additional
 L-glutamine is added to the HBM medium of the present
 invention as described to compensate for this.

10 Table III below lists substances which may be
 optionally added to HBM medium to optimize cell growth for
 specific purposes and for specific species origin of
 hepatocytes and specific cell lines or malignant cells.
 Again, these amounts will vary depending on whether a
 particular component is already present in the stock basal
 medium being used.

15 TABLE III
Optional Additives for HBM

		<u>Preferred Amount</u>	<u>Concentration Range</u>
		<u>units/L</u>	<u>units/L</u>
20	Albumin*	2.0 g	0-10.0 g
	Penicillin**	100 U	0-2500 U
	Streptomycin**	100 µg	0-2500 µg
	Sodium Pyruvate	0.15 g	0-2.0 g
	Gamma Tocopherol	0.35 mg	0-3.5 mg
25	Alpha Tocopherol	0.15 mg	0-2.5 mg
	Vitamin D ₃	0.20 mg	0-1.8 mg
	Dextran*	2.0 g	0-5.0 g
	Iron Gluconate***	25.0 µg	0-100 µg
	Linolenic Acid	1.0 g	0-5.0 g
30	Apo-transferrin***	5.0 mg	0-20.0 mg
	Retinol	0.05 mg	0-2.0 mg
	Vitamin B ₁₂	0.15 mg	0-2.0 mg
	Ascorbic acid	3.0 mg	0-10.0 mg
	Choline Chloride	1.5 mg	0.2-12.5 mg
35	Biotin	0.75 mg	0.2-15.0 mg

- * Dextran can be substituted for albumin.
- ** Other antibiotics may be used, such as for example, gentamycin.
- 5 *** Iron gluconate and apo-transferrin can be substituted for iron-saturated holo-transferrin.

In instances where accelerated growth is desired various growth factors can be added to the HBM of the present invention. While HGF/SF, EGF, and TGF α are
10 presently preferred, it is to be understood that other growth factors may also be used. The preferred amount of such growth factors typically vary with the particular source used. The amounts listed herein are therefore necessarily limited to the particular sources used herein.
15 In the Example below HGF/SF was preferably added to HBM at 40 ng/ml; EGF was preferably added at 20 ng/ml; and TGF α was preferably added at 20 ng/ml.

The following examples are included for illustrative purposes only and are not intended to limit
20 the scope of the invention.

EXAMPLE

Materials and Methods

Materials

Male Fischer 344 rats from Charles River,
25 (Pennsylvania) were used for all the experiments involving rat hepatocyte isolation. EGF and Matrigel (a mixture of matrix components derived from EHS mouse tumor) were obtained from Collaborative Research (Waltham, MA).
[³H]Thymidine was obtained from ICN Radiochemicals
30 (Irvine, CA). Collagenase for hepatocyte isolation was obtained from Boehringer Mannheim (Indiannapolis, IN).

Vitrogen (Celtrix Labs, Palo Alto, CA) was used for the construction of the collagen gels. General reagents were obtained from Sigma Chemical Co. (St. Louis, MO). HGF/SF
5 used for these studies was the Δ5 variant. ECL matrix was purchased from Upstate Biotechnology (Lake Placid, NY).

Isolation and Culture of Hepatocytes

Rat hepatocytes were isolated by an adaptation of the calcium two step collagenase perfusion technique as
10 taught by Kost, D.P., et al, J. Cell. Physiol. 147:274 (1991), the disclosure of which is incorporated herein by reference. All such preparations were designed to obtain a population of pure hepatocytes. Unlike many other hepatocyte culture systems, the present invention does not
15 require or use a feeder cell co-culture or feeder-cell conditioned media. After isolation of hepatocytes, the cells were suspended in a media used for cell attachment to the culture plates. This media was MEM (GIBCO 12570) with NEAA (GIBCO 11140), insulin 5 mg/L, and gentamycin
20 5 µg/ml. Hepatocytes were plated on a single layer of collagen coating as described below and left to attach for 2 hours. Six-well cluster plates (9.8 square centimeters per plate) from Corning were used. For experiments where hepatocytes were not going to be induced to proliferate,
25 or where assessments of differentiated function were performed, the cells were plated at a density of 80,000 hepatocytes per square centimeter of surface area. For experiments intended to induce proliferation or genetic transduction with foreign DNA, cells were plated at an
30 initial density of 1,000 or 10,000 hepatocytes per square centimeter of surface area. The plating medium was replaced with the invention HBM medium at 2 hours after the cells were plated and every 48 hours thereafter. Thymidine, growth factors, and other ingredients, were
35 added at the time of medium change as required.

Human hepatocytes were isolated by an adaptation of the collagenase perfusion technique as described by Strom, S.C., et al., Journal of National Cancer Institute 65(5):771-8 (1982), the disclosure of which is incorporated herein by reference. Cells were cultured as described above for the rat hepatocytes.

Collagen gels were prepared as described by Michalopoulos, G.K., et al., Exp. Cell. Res. 94:70 (1975), the disclosure of which is incorporated herein by reference. Dry coating of plates with collagen and Matrigel was also performed as specified by the manufacturer. Matrigel gels were made by adding 50 μ l of Matrigel solution into 0.5 ml of medium directly on the top of attached cells.

DNA synthesis was measured by the uptake of tritiated thymidine into trichloroacetic acid (TCA) precipitable material as described by Kost, D.P., et al. (1991), cited above. Collagen gels, where necessary, were digested with 2 mg of collagenase per ml of MEM medium used. Incubation was then carried out for 30 minutes at 37°C. The digested gels were treated with NaOH followed by TCA to precipitate DNA, RNA, and proteins as described by Kost, D.P., et al. (1991), cited above.

Composition of the HBM Medium

DMEM, HEPES, L-glutamine, and antibiotics were purchased from GIBCO/BRL (Gaithersburg, MD). ITS mixture (Insulin, Transferrin, Selenium) was purchased from Boehringer Mannheim. All other additives were cell culture grade (Sigma). Unless otherwise indicated for specific experiments, the stock basal medium consisted of DMEM 11966 and DMEM 10314 blended to achieve a final D-glucose concentration of 2.0 g/L. In this case, 445 ml of DMEM 10314 was blended with 555 ml of DMEM 11966 to

achieve this concentration. The formulas for these stock basal media are set forth in Table I above. The resulting blended medium was then supplemented with: purified
5 bovine albumin 2.0 g/L, D-galactose 2.0 g/L,
L-ornithine 0.1 g/L, L-proline 0.030 g/L, nicotinamide
0.610 g/L, ZnCl₂ 0.544 mg/L, ZnSO₄·7H₂O 0.750 mg/L,
CuSO₄·5H₂O 0.20 mg/L, MnSO₄ 0.025 mg/L, glutamine 5.0 mM,
ITS (rh-insulin 5.0 mg/L, human transferrin 5.0 mg/L [30%
10 diferric iron saturated], selenium 5.0 µg/L),
dexamethasone 10⁻⁷ M, and HEPES buffer 20.0 mM.
Penicillin and streptomycin were added at 100 U/L and
100 µg/L, respectively. The mixed basal HBM was
sterilized by filtration through a 0.22-µm low protein-
15 binding filter system (Corning), stored at 4°C, and used
within 4 weeks. The growth factors, as required, were
added to HBM fresh at the specified concentrations every
time the medium was changed.

Retroviral Transfection and Assessment of Clonal Expansion

20 Hepatocytes were initially plated at 10⁴/cm² and
grown in HBM supplemented with HGF/SF (40 ng/ml) and EGF
(20 ng/ml). After 68 hours the media was replaced with
supernatant from CR ψ P-packaged, replication-deficient,
amphotropic retrovirus (MFG - 5 x 10⁵ units per ml)
25 containing the *E. coli* β -galactosidase gene under an LTR
promoter as described by Zitvogel, L.H., et al., Hum.
Gene. Ther. 5:1493 (1994), the disclosure of which is
incorporated herein by reference. Polybrene was added at
2 µg/ml. The supernatant was replaced after 18 hours with
30 HBM that was supplemented with EGF at 20 ng/ml and HGF/SF
at 40 ng/ml. The brief exposure with the virus-containing
supernatant did not have an adverse effect on hepatocyte
survival or proliferation. At indicated times, cells were
fixed with 0.5% glutaraldehyde in PBS for 10 min and
35 developed with X-Gal substrate at 37°C for 16 hours.

Transduced cells expressing the *E. coli* gene stained positive as shown in Figures 4G and 4F. Appropriate controls for each component were negative for X-Gal staining.

Transmission Electron Microscopy

Samples for transmission electron microscopy (TEM) were fixed on the culture plates for 1-1.5 hours in 0.1 M sodium cacodylate buffer (pH 7.4) that contained 2.5% glutaraldehyde and 2% formaldehyde. The plates were then rinsed 2 times with 0.1 M sodium cacodylate buffer (pH 7.4) and 2 times with 0.1 M sodium cacodylate buffer containing 5% sucrose (pH 7.4). They were held in the sucrose buffer for 1-7 days, rinsed 2 times with 0.1 M sodium cacodylate buffer (pH 7.4), and then postfixed for 1 hour in 1% OsO₄ in 0.1 M sodium cacodylate buffer. The plates were then rinsed again in buffer, and the fixed and processed collagen gels were then cut in strips with a razor blade. The strips were then transferred to glass specimen vials, dehydrated through a graded series of ethanol (25-100%) and two propylene oxide changes, and infiltrated with Epon-Araldite resin (BioTec, TX). Several changes of resin were made over 2 days, as the collagen gels tended to hold the propylene oxide. The collagen strips were flat-embedded and cured overnight at 60°C.

Analysis of Gene Expression by Northern Blots

Extraction of Total RNA and mRNA from Cultures.

Total RNA was extracted from unwashed cell cultures using 2.0 ml of RNAzol B (BioTec) per well and purified per the manufacturer's guidelines. RNA concentration and purity were determined by routine spectrophotometry. Size separation of 20 µg RNA per lane was completed on denaturing 1% agarose gels and transferred to nylon membranes (Amersham, Arlington Heights, IL) by the

capillary method. After cross-linking under UV light, the membranes were hybridized overnight with specific cDNAs (as indicated in the figures) that had been labeled with [α-32P]dCTP using an Amersham random primer kit. Membranes were subsequently washed under high stringency conditions and exposed to XAR film (Eastman Kodak, Rochester, NY) for 1-3 days. Quantification of the RNA hybridization bands were performed by laser densitometry.

10 Sources of cDNA Probes

cDNA probes used to study gene expression were obtained as gifts, and are available upon request from the following sources: Cytokeratin 8 from Dr. Norman Marceau (Laval University); Cytokeratin 14 from Dr. Dennis Roop (Baylor College of Medicine); Cytokeratin 18 from Dr. Robert Oshima (LaJolla Cancer Research Foundation); Cytokeratin 19 from Dr. Andre Royal (University of Montreal); TGFα (rat) originated from Dr. David Lee (University of North Carolina at Chapel Hill); EGF-R (rat) originated from Dr. Sheldon Earp (University of North Carolina at Chapel Hill); aFGF from American Type Culture Collection (ATCC) (catalog No. 78222); aFGF-R from ATCC (catalog No. 65796); uPA originated from Dr. Jay Degen (University of Cincinnati); cytochrome 11B1 from Dr. Steve Strom (University of Pittsburgh); cDNAs for albumin, α fetoprotein, and transcription factor analysis were generated by Dr. Joe Locker (University of Pittsburgh).

RESULTS OBTAINED

30 Role of Media Components and Matrix Substrates on Cell Proliferation

The full description of the HBM medium is given above. To evaluate the relative importance of different media components several experiments were performed whose results are shown in Table IV (A, B, and C) below. The

components D-glucose, albumin, dexamethasone, transferrin and selenium, nicotinamide, and trace elements were individually subtracted from the full HBM medium composition as shown in Table IV A. The total DNA per culture after 14 days of growth is shown therein. As can be readily seen, removal of dexamethasone had the most dramatic effect, followed by removal of nicotinamide. In Table VI B, growth of cells achieved by day 14 is compared between HBM medium containing diferric transferrin (iron saturated) versus iron unsaturated transferrin. Addition of iron containing diferric transferrin (30% saturation) was found to be much more effective in promoting growth. The addition of elemental iron (FeSO_4 , $0.1 \mu\text{M}$) to the unsaturated transferrin failed to overcome the difference. Table VI C provides information on the relative effects of D-glucose, D-galactose, and L-ornithine in HBM medium. All three of these components are potential sources of energy to the cells. Complete cessation of growth was noticed when all three components were removed. Addition of D-glucose alone restored most of the response whereas the addition of D-galactose alone was less effective. Ornithine alone had minimal effect. Concentrations of 2 g per liter each of albumin and D-glucose were found to be optimal though the effects were not statistically different than 1 or 3 g per liter in each case. The effect of complete removal of these components is shown in Table IV A.

Table IV

A. Effect of Removal of Specific HBM Components on Hepatocyte Growth

	<u>µg/well</u>
5	Zero time DNA 13.90 ± 3.60
	DNA at day 14 in HBM (with HGF/SF and EGF) supplemented with:
	+ All components 84.90 ± 0.50
10	- Glucose 69.80 ± 2.90
	- Albumin 68.70 ± 0.50
	- Dexamethasone 13.70 ± 0.20
	- (Transferrin and selenium) 63.10 ± 2.00
	- Nicotinamide 35.20 ± 1.70
15	- Trace elements 65.00 ± 2.40
	- All components 20.20 ± 4.30

B. Effects of Iron and Transferrin on Hepatocyte Growth

	Zero time DNA 9.21 ± 1.01
	Diferric transferrin (iron saturated) 70.15 ± 1.21
20	Diferric transferrin plus added iron (5 µM) 1.83 ± 0.41
	Iron poor transferrin 24.84 ± 4.30
25	Iron poor transferrin with added iron 1.10 ± 0.90

C. Effect of Removal of Glucose, Ornithine, or Galactose

	Zero time 11.0 ± 0.4
	Control (Gluc.+, Orn.+, Gal+) 59.0 ± 2.5
	Gluc.-, Orn.-, Gal.- 10.6 ± 0.8
30	Gluc.-, Orn.+, Gal.- 14.4 ± 0.9
	Gluc.-, Orn.-, Gal.+ 44.2 ± 6.0
	Gluc.+, Orn.-, Gal.- 59.0 ± 4.0
	Gluc.+, Orn.+, Gal.- 62.7 ± 1.4
	Gluc.+, Orn.-, Gal.+ 63.6 ± 0.3
35	The indicated HBM components were removed and hepatocyte were grown in the modified media for 14 days. The total DNA per culture in micrograms was measured at day 14 to evaluate growth cell growth. Time zero was the sample of hepatocyte suspension that was inoculated into the plate immediately after cell isolation. The data are expressed as mean ± standard error of three separate plates.
40	

Hepatocytes Enter Diffusely into Proliferation under the Influence of HGF/SF, EGF, and TGF α

5 Figures 1A and 1B show the uptake of thymidine
per μ g DNA, as well as the BRdU nuclear labeling index at
different days in culture in cells growing in the presence
of HGF/SF and EGF (40, 20 ng/ml respectively) as described
above. As can be seen, most of the proliferation occurs
at days 5-12. By day 15 the cultures were confluent and
10 DNA synthesis slowed down. The high nuclear labeling
index during the times of sustained proliferation
indicates that the proliferating cells derive directly
from the mature hepatocytes.

15 The growth factors HGF/SF, EGF, TGF α , KGF
(keratinocyte growth factor), SCF (stem cell factor), and
aFGF (acidic fibroblast growth factor) were added
individually to HBM medium. Of the growth factors added,
HGF/SF, EGF, and TGF α (shown in Figure 2 as "TGFA") caused
significant cell proliferation, as shown by the total
20 amount of DNA per culture at day 15. KGF, aFGF, and SCF
when added alone or in combination had no proliferative
effect as shown in Figure 2. TGF α had a stronger
proliferative effect than any of the other mitogens when
added alone to HBM medium. HGF/SF and EGF together had
25 the strongest proliferative effect for a given time
interval of 15 days together than did any other single
mitogens or combinations. Addition of all the growth
factors together had no more effect than the combined HGF/
SF and EGF as seen in Figure 2. The detailed cell
30 kinetics induced by HGF/SF and EGF alone or in combination
are shown in Figure 1C. The total DNA per culture is
shown as a function of the time in culture. The largest
amount of accumulated DNA at day 15 was seen with the
combination of HGF/SF and EGF. The DNA per culture at day

15 was 12 times that at time 0, reflecting the increase in cell number. HGF/SF and EGF were about equally potent. TGF α alone was more mitogenic than either HGF/SF or EGF alone.

Effect of Matrix Substrates

Several matrix substrates promoted cell growth in this system. Rat hepatocytes were cultured for 15 days in HBM supplemented with HGF/SF (40 ng/ml) and EGF (20 ng/ml) grown on various matrices as seen in Figure 3. The cells were cultured as described above. Dry coating with collagens type IV (mouse), type I (bovine), fibronectin, and laminin were equally effective in promoting cell growth as assayed by measurement of total DNA per culture at day 15. Dry coating with ECL (a commercial derivative of EHS gel, UBI) had superior effects. Coating with type I collagen (Vitrogen commercial preparation) was the standard method used for the experiments unless otherwise specified. The effect of matrix gels in promoting specific phenotypic conversions in these cultures is further discussed below.

Phenotypic Changes of Hepatocytes During Proliferation

The morphology of the proliferating cells (cultured in HBM as described above supplemented with 40 ng/ml HGF/SF and 20 ng/ml EGF) varied at different times after the stimulation of cell proliferation. From a normal hepatocyte morphology as seen in Figure 4A, the proliferating cells in the first 4 days acquired long projections assuming the phenotype typically described as due to the "scattering" effect of HGF/SF on hepatocytes as seen in Figure 4B as described by Michalopoulos, G.K., et al, J. Cell. Physiol. 156:443 (1993), the disclosure of which is incorporated herein by reference. Between days 6 and 8, the proliferating cells lost most of their

cytoplasmic granules, the nuclei became less prominent, the projections diminished and the cells began to grow as monolayer patches. Eventually these patches merged as the cells continued to grow to form a continuous monolayer as seen in Figure 4C. Examination by electron microscopy seen in Figures 4D and 4E showed that most of the features typical of mature hepatocytes were missing. By day 15 there were no lamallae of endoplasmic reticulum wrapping around mitochondria and there are no glycogen rosettes or peroxisomes. Bile canaliculi were absent. There was a prominent increase in bundles of keratin intermediate filaments. The nuclei were angular with very prominent nucleoli. After confluency the morphology gradually revert back to mature differentiated hepatocytes with endoplasmic reticulum lamallae, mitochondria, glycogen, peroxisomes, bile canaliculi, etc., similar to that in Figures 6B and 6C.

Clonal growth of the proliferating rat hepatocytes is demonstrated in Figures 4F and 4G. Hepatocytes were transfected at day three in culture with a replication-deficient retrovirus containing the lac-Z gene under the influence of a viral LTR and were stained for expression of β -galactosidase. Mostly single cells were stained positive at day four in culture (1 day after transfection) as seen in Figure 4F. On the other hand, staining at day 10 and continuing through day 28 showed patches of positive-stained, hepatocytes, consistent with clonal growth of the original transfected hepatocytes as seen in Figure 4G. The percentage of lac-Z-positive cells (~20%) did not appear to change during culture.

Proliferating Hepatocytes Express Altered Levels of Various Genes

The expression of several specific genes was assayed in proliferating hepatocytes. These included mRNA genes associated with hepatocyte differentiation (albumin,

cytochrome IIB1 (labeled as P450 in Figure 5A)), genes encoding cytokeratin markers (cytokeratins 14, 18, and 19) or related to hepatocyte growth (urokinase (uPA), HGF/SF and its receptor c-met (labeled as MET in Figure 5A), EGF (labeled as EGFR) and TGF α and their receptor, acidic FGF and its receptor, and TGF β 1). These genes were studied by Northern blot analysis of RNA from cultures grown in the presence of either combined HGF/SF (40 ng/ml) and EGF (20 ng/ml) (Figures 5A and 5B) or with TGF α alone (20 ng/ml) (data not shown). Total RNA was isolated from cultures at days 0, 6, 10, 15, and 21. No expression of HGF/SF or TGF β 1 mRNA was seen at any of the time points examined.

As can be seen, albumin and cytochrome IIB1 mRNA were present at time zero and subsequently decreased. Albumin mRNA increased by day 21 at which time mRNA for α fetal protein (AFP) was also detected. mRNA for cytokeratins, 14, 18, and 19 increased through culture. There was a steady increase in mRNA for aFGF and TGF α . The mRNA of receptors for HGF/SF (MET, Figure 5A) and aFGF (Figure 5B aFGFR) remained present throughout the culture time. The EGF receptor (EGFR) mRNA declined from day zero but remained expressed. GAPDH mRNA expression was used as a reference "housekeeping" gene. Dramatic increases were noted in expression of urokinase as well as cytokeratins 14 and 19. Some differences from the above pattern were seen in the cultures that, instead of HGF/SF and EGF, were maintained in the presence of TGF α (20.0 ng/ml). In these cultures albumin expression and expression of the HGF/SF receptor were better preserved during proliferation whereas AFP appeared earlier. (Data not shown.) Despite the observed differences in gene expression patterns no morphologic differences were seen between cells growing in the presence of TGF α or HGF/SF plus EGF, once confluency was achieved.

Proliferating Hepatocytes Revert to Mature Hepatocytes
under the Influence of Matrigel or in the Presence
Nonparenchymal Cells or With Time in Culture

5 When Matrigel was overlaid on cultures at day 8,
there was a rapid (within 2 days) appearance of bile
canaliculi and organization of the cells into cord-like
structures. The features of these cells are shown in
Figure 6A. As shown by electron microscopy in Figures 6B
10 and 6C, these cells had typical markers of mature
hepatocytes, including wrapping of the endoplasmic
reticulum around mitochondria, bile canaliculi, and the
presence of glycogen. Preparations of mRNA were made from
cultures exposed to Matrigel for 10 days (days 8-18 in
15 culture). The expression of albumin was compared between
day zero in culture (immediately after collagenase
perfusion), at day 8 in culture (before the overlay by
Matrigel), and cultures at day 3 and 7 after Matrigel
overlay as shown in Figure 6D. Addition of Matrigel
20 caused dramatic increases in expression of albumin mRNA,
compared to control proliferating cultures in which it was
minimally detectable. The effect of Phenobarbitol (PB) on
the levels of cytochrome P450 IIB1 mRNA in the Matrigel-
treated cultures was also measured as shown in Figure 6E.
25 Matrigel was added to the cultures at day 8. PB was added
2 days later (day 10 of culture). The cells were
harvested 5 days after addition of PB (day 15 in culture).
Addition of PB induced cytochrome IIB1 mRNA only in the
Matrigel-treated cultures. Induction of this mRNA by
30 phenobarbitol is typical of hepatocytes and does not occur
in any other cell as reported by Michalopoulos, G., et
al., Science (Wash. DC) 193:907 (1976), the disclosure of
which is incorporated herein by reference. Typically,
hepatocyte cultures rapidly lose the capacity to respond
35 to PB. This finding is supporting evidence that addition
of Matrigel to the cultures of proliferating hepatocytes
induces a mature hepatocyte phenotype, as attested to by
the electron microscopic structure shown in Figures 6B and

6C. The expression of cytokeratin 19 (CK19) (Figure 6F),
a bile duct marker expressed by the proliferating
hepatocytes before introducing differentiating conditions,
5 also ceased expression after addition of Matrigel as seen
in Figure 5B.

DNA synthesis was measured in the cultures
exposed to Matrigel and there was a substantial decrease.
To assess whether differentiation to mature hepatocyte
10 morphology required DNA synthesis, 20 mM hydroxyurea was
added to the HBM media. This has been shown
(Michalopoulos, G.K., et al., Cancer Res. 38:1866 (1978),
the disclosure of which is incorporated herein by
reference) to abolish scheduled semiconservative DNA
15 synthesis in hepatocytes by inhibiting ribonucleotide
reductase. Hydroxyurea was added to cultures before the
Matrigel overlay and maintained throughout the next 5-day
period. DNA synthesis decreased down to 3.93% of the
control (without hydroxyurea) in the proliferating
20 cultures maintained in the absence of Matrigel and down to
6.27% of control (without hydroxyurea) in the cultures
maintained in the presence of Matrigel. Though DNA
synthesis was decreased down to 6.27% of control
(+Matrigel, no hydroxyurea) levels, the conversion of the
25 proliferating hepatocytes to mature hepatocyte morphology
was entirely unaffected and involved the entire
population.

HGF/SF (but Not TGF α or EGF) Induces Proliferating
Hepatocytes to Differentiate into Ductular/Acinar
30 Structures in Type I Collagen Gels

Hepatocytes maintained between two collagen gel
layers retain their morphology and differentiation for
prolonged time periods as described by Michalopoulos,
G.K., et al., J. Cell. Physiol. 156:443 (1993), the
35 disclosure of which is incorporated herein by reference.

Hepatocytes maintained in collagen gel sandwiches in cultures with conventional media containing HGF/SF undergo intense proliferation, form prominent projections, and eventually become organized in structures reminiscent of the hepatic plates. The behavior of hepatocytes maintained between two layers of collagen gel as previously described was examined, but in the presence of HBM supplemented with either HGF/SF or EGF as described above. It was noted that in the EGF supplemented media hepatocytes underwent the typical phenotypic transitions as described above. On the other hand, in hepatocytes in HBM supplemented with HGF/SF alone, following expansion of cells identical in appearance to the proliferating hepatocytes described above, there appeared multiple duct-shaped structures between days 10 and 15. These became prominent and encompassed most of the cells present in the cultures. Starting from approximately day 10 and by day 15 most of the cells in the culture were arranged in such ductular structures. The appearance of these structures is shown in Figure 7A. Histologic sections are shown in Figure 7B (light microscopy) and Figures 7C and 7D (electron microscopy). The structures had a ductular or acinar configuration. Some of the cells surrounding these structures were very attenuated and had light and electron microscopic appearance identical to bile duct epithelium. Others however are larger and resemble more closely the ductular hepatocytes described in previous studies of in vivo models. The proliferation of cells (data not shown) under either EGF or HGF/SF in the collagen gel sandwiches was much less (<25% at the highest peak) than that seen in the cultures on plastic coated with dry collagen. Most proliferation ceased by day 10 and the duct-like structures appeared after cell proliferation had ceased (days 10-15). Ductular acinar structures were also noted in these cultures when HGF/SF and EGF were combined but were fewer than with HGF/SF alone. As with the Matrigel overlay, additional of hydroxyurea to inhibit DNA

synthesis (inhibition down to 5.1% of control) did not affect the formation of the ductular structures (data not shown). Figure 7E shows that these cells express
5 cytokeratin 19 which is characteristic of duct cells. (See, Sirica, A.E., Prog. Liver Dis. 10:63 (1992) and Sirica, A.E., Histol. Histopathol. 10:433 (1995), the disclosures of which are incorporated herein by reference.) A small amount of albumin expression was also
10 retained, consistent with the presence of hepatocyte-like cells within the ductular structures as seen in Figure 7D. It was noted that the ductular cells maintain CK19 expression in the nonproliferating state (Figure 7F) in contrast to the cells differentiating toward the mature
15 hepatocyte lineage which cease expressing this bile duct marker.

Sustained Growth and Population Expansion of Human Hepatocytes in HGM, in the Presence of HGF/SF and EGF

Though human hepatocytes cultures have not been
20 characterized as extensively as those of the rat, the literature available has shown that these cells also undergo a limited round of DNA synthesis after stimulation by growth factors and rapidly degenerate in culture. See, Ismail, T., et al., Hepatology 14:1076 (1991), the
25 disclosure of which is incorporated herein by reference. The response of human hepatocytes to HGF/SF (40 ng/ml) and EGF (20 ng/ml) in HBM medium was studied. Results similar to those described for rat hepatocytes were found in
30 primary cultures of human hepatocytes, as seen in Figure 8. The human cells begin to rapidly proliferate at day 3-4 in culture and reach confluency by day 19.

As stated above, there is a present need for a media that allows hepatocytes that are cultured in vitro to expand as a cell population. The HBM of the present
35 invention allows such expansion and therefore enables much

needed research of proliferating hepatocytes. The present invention also will be useful in numerous other applications outlined below.

5 For example, all current methods in liver-targeted gene therapy that achieve stable, long-term expression of transferred genes require actively dividing cells during the initial transfection. Normal liver has only 1 in 20,000 hepatocytes in S phase growth on any particular day. In order to increase cell proliferation, a large portion (2/3) of the patient's liver must be removed. Subsequently the liver-targeted gene carrier source is intravascularly injected. The alternative to this drastic measure is to remove a small piece of liver (10%), culture the hepatocytes, transfect them in culture, and then reinfuse the cells back into the liver. This latter method, while safer, less expensive and better controlled than the former, is currently hampered by the lack of culture media such as HBM which allows prolonged proliferation, clonal expansion of essentially all cells cultured, and long term viability of the cells.

 The design of current extracorporeal liver devices based on bioartificial liver tissue which include cellular elements rely on transformed (tumor) hepatocytes or animal hepatocytes. The animal cells do not survive for more than few days in the bioreactors that are used and do not proliferate, making it necessary to generate bioreactors more frequently and in close proximity and to maintain all necessary facilities, materials and animal donors in anticipation of need. Additionally, animal cells may have several undesirable aspects which makes them clinically difficult or dangerous to use. Also, tumor cells used in these devices carry the risk of leakage of cells into the patient and therefore potentially could give rise to tumors in the patient. The HBM media of the present invention will permit production

of bioreactors containing animal and/or human hepatocytes and/or other cells and provide the unique features of increasing cell numbers while maintaining long term viability and complete differentiation. Such reactors would also be useful for drug manufacture, drug metabolism, toxicology studies, and complex biological studies.

Another potential use for HBM media of the present invention is in autotransplantation of hepatocytes. Hepatocyte transplant is a relatively new approach to treating patients with endstage liver disease. The method relies on use of donor organs which are not used for organ transplant. Autotransplantation has the advantage of not requiring immunosuppressive drugs, but it is not practical if it is not possible to expand the number of cells and keep them alive long enough to place them back into the donor. Endstage liver explants could also provide a donor hepatocyte source under selected conditions, including encapsulation or genetic manipulation. Regardless of the source of hepatocytes, expansion of cell number in long term culture and replantation (into spleen, portal system, peritonium, renal capsule, etc.), does enhance effective hepatic synthetic and detoxification processes for prolonged periods of time. The limiting point is the lack of hepatocytes. HBM has the potential to eliminate this lack of cells, and thereby dramatically expand the number of patients that could be treated for liver failure. Similarly, autotransplantation or heterotransplantation with other cells grown in the media of the present invention could expand the number of patients and array of treatments possible with biocellular therapies.

The HBM of the present invention can also be useful for drug and chemical testing procedures. Virtually all drugs, chemicals, and other manufactured products must be tested for mutagenicity and toxicity in hepatocyte cultures as part of FDA, USDA, NIOSH and EPA regulations. Currently, these tests are performed on rat hepatocytes in short term culture because of lack of long term viability or significant proliferation. The use of HBM will (1) extend the testing period (longer term viability), (2) allow longer exposure periods at lower concentrations, (3) allow observation of mutagenicity and toxicity on proliferating hepatocytes, and (4) make available a supply of human hepatocytes which could be used in other toxicologic investigations.

Although the invention has been described in detail for the purpose of illustration, it is to be understood that such detail is solely for that purpose and that variations can be made therein by those skilled in the art without departing from the spirit and scope of the invention except as it may be limited by the claims.